

DIRECTIONS FOR USE

AGROBOR™ is recommended for use on all crops during plant growth periods that require relatively high levels of Boron. AGROBOR™ is a fully water soluble, effective, readily available source of Boron for soil and foliar applications.

AGROBOR™ will be influenced by differences in crop varieties, soil and environmental conditions and should be applied under conditions of optimum plant growth. Optimum conditions are defined as adequate humidity, moderate temperatures and adequate soil moisture. Optimum conditions for foliar application to most crops generally occur during early morning or late evening hours.

- Do not apply to crops that are undergoing a period of moisture or heat stress or in very windy conditions.
- Wear suitable protective clothing when sprayed with other products.
- For optimum results with this product, the pH of the tank mixture should be between 5 and 6.
- This product should be applied as a foliar spray when crops have sufficient leaf area.
- For optimum results AGROBOR™ should be used in a plant nutrition program. Recommendations should be based on soil and leaf analyses.
- Use the lower dosage per 100 gal at higher water volumes and higher dosage per 100 gal at lower spray volumes.
- Where overhead systems are in use, avoid scheduling irrigation directly after application.

- In orchards equipped with micro irrigation systems, spray deposits should not be affected by irrigation.

CAUTION

Do not apply on wet fruit. Application can only take place once the fruit is dry. Do not apply to plants that are undergoing a period of moisture or heat stress, or during cool wet conditions. Do not spray AGROBOR™ within 30 days after application of any Copper containing products. Do not store mixed product for more than 8 hours. Do NOT mix or apply in combination with Calcium containing products, amine herbicides, Copper containing fungicides, Sulfates, Phosphates, phosphonates, highly alkaline material such as Lime Sulfur, MCPA or highly alkaline Boron products. Do not exceed the appropriate application rates.

CAUTION

AGROBOR™ contains Boron. Use of AGROBOR™ on any crops other than those recommended may result in serious injury to the crop(s).

Consult with your distributor, crop adviser or manufacturer should crop specific programs, or any other information be required around the use of this product.

STORAGE & DISPOSAL

Store product in original container. DO NOT store in excessive cold or heat or direct sunlight. DO NOT reuse container. DO NOT contaminate water, food or feed by storage or disposal. Keep packaging closed and securely fastened.

AGROBOR™		F002806
		
Agri Technovation Inc. 516 Villa Ave, Ste 23, Clovis, CA, 93612		
		
GUARANTEED ANALYSIS		
Boron (B)	16.00%	
Derived From: Boric Acid		
WATER-SOLUBLE CRYSTALLINE BORON COMPOUND Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.html		
KEEP OUT OF REACH OF CHILDREN. DANGER May damage fertility or the unborn child.		
 Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Wear protective gloves, protective clothing and eye protection. IF exposed or concerned: Get medical advice. Store locked up. Dispose of container to in accordance with regulations.		
NET WEIGHT: 50 LBS (22.68 kg)		

APPLICATION RATES

The rate of application is dependent on the soil fertility and reaction required. For more information on foliar application rates and rates for applying through the irrigation system, see table on application rates. For other modes of application, consult a qualified advisor. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies.

MIXING

Always mix AGROBOR™ with sufficient quantities of water. Never apply AGROBOR™ undiluted. Follow standard tank mixing procedures. Ensure good agitation throughout application.

DISCLAIMER

The manufacturer, importer and seller make no express or implied warranties, including any warranty of merchantability, fitness for a particular purpose, or suitability. The product is sold "AS-IS". All sales are final. In no event will manufacturer, importer and seller have any liability for any damages, whether direct, incidental, special or consequential arising out of or in connection with any duty with respect to the product, including, but not limited to lost profits, lost sales or injury to persons or property. Manufacturer, importer and seller's liability on any claim of any kind, including but not limited to warranty, negligence, strict liability, and any other cause of action, for any loss or damage arising out of, connected with, or resulting from the performance or breach of the terms of sale of any goods, or any related services related to sale, delivery, resale, technical direction, inspection, operation or use of any goods or services or part of those goods or services will in no case exceed the purchase price allocable to the goods or services or part of those goods or services that gives rise to the claim. Manufacturer, importer and seller make no representation whatsoever with regard to whether any goods comply with the Occupational Safety and Health Act (OSHA), the Americans with Disabilities Act (ADA), or any other federal, state or local statute, law, ordinance or ruling. Customer, buyer and user assume all responsibility for use of product.

COMPATIBILITY

It is advisable to do a miscibility test prior to mixing with any other product and to consult with a qualified advisor or distributor.

APPLICATION RATES

Crop	Apply as Foliar Application or Through Irrigation	Remarks
Cabbage Crops	1 – 2 lbs/acre through spraying in 50 – 80 gal of water or 3 – 4 lbs/acre through irrigation	For foliar application, apply as a full cover spray, and repeat only when you have consulted with your agronomist or horticulturist.
Carrots	1 – 2 lbs/acre through spraying in 50 - 80 gal of water or 2 - 3 lbs/acre through irrigation	For foliar application, apply as a full cover spray, and repeat only when you have consulted with your agronomist or horticulturist.
Citrus	Foliar application 1 lbs/100 gal of water.	First spray - Apply before 10% flowering. Second spray – Apply at 100% petal fall. Do not apply to young trees. Apply as a light-cover spray.
Cotton	1 lbs/acre through spraying or 2 - 4 lbs/ acre through irrigation	For foliar application, apply as a full cover spray, and repeat only when you have consulted with your agronomist or horticulturist.
Deciduous Fruit	Foliar application 1 lbs/100 gal of water.	Apply as a light-cover spray at 30% blossoming and again after harvest in autumn. Do not exceed a 0,1% concentration.
Flowers and Ornamentals	Foliar application 1 lbs/100 gal of water.	Apply as a full cover spray, and repeat only when you have consulted with your agronomist or horticulturist.

APPLICATION RATES

Crop	Apply as Foliar Application or Through Irrigation	Remarks
Grapes	Foliar application 2 lbs/100 gal of water	Apply just before flowering and again directly after harvest. Do not exceed a 1% concentration.
Alfalfa	1 – 2 lbs/acre through spraying or 4 lbs/gal through irrigation	When applied through spray tank apply as a full cover spray, and repeat only when you have consulted with your agronomist or horticulturist.
Corn	1 – 2 lbs/acre through spraying or 3 - 4 lbs/ acre through irrigation	Apply 4 weeks after emergence and 2 weeks before silking. Consult with your agronomist about the maximum pure Boron per acre, per crop.
Nuts	Foliar application 1 lbs/acre of water.	First spray - Apply before 10% flowering. Second spray – Apply at 100% petal fall. Do not exceed the maximum rate. Apply in 200 – 300 gal/acre as a light-cover spray.
Onions	1 – 2 lbs/acre through spraying in 50 – 80 gal of water or 3 – 4lbs/acre through irrigation	Apply as a full cover spray and repeat only when you have consulted with your agronomist or horticulturist.
Potatoes	1 – 2 lbs/acre through spraying in 50 – 80 gal of water or 3 – 4 lbs/ acre through irrigation	Apply 1 week after initiation and repeat thereafter, only when you have consulted with your agronomist or horticulturist.

APPLICATION RATES

Crop	Apply as Foliar Application or Through Irrigation	Remarks
Strawberries	1 – 2 lbs/acre through spraying in 50 – 80 gal of water or 2 – 3 lbs/acre through irrigation	Apply at 8 - 10 leaf stage and repeat only when you have consulted with your agronomist or horticulturist.
Sugar Beet	1 – 2 lbs/acre through spraying in 50 – 80 gal of water or 3 – 4 lbs/acre through irrigation	Apply as a full cover spray and repeat only when you have consulted with your agronomist or horticulturist.
Subtropical Fruit	Foliar application 1 lbs/ 100 gal.	First spray -Apply before 10% flowering. Second spray – Apply at 100% petal fall. Do not exceed the maximum concentration. Apply in 200 – 300 gal as a light cover spray.
Sugarcane	1 – 2 lbs/acre through tractor spray or 3 – 4 lbs /acre through irrigation	Apply 4 weeks after emergence and 2 weeks after pluming. Consult with your agronomist about the maximum pure Boron per acre, per crop.
Sunflower	1 – 2 lbs/acre through tractor spray or 3 – 4 lbs/ acre through irrigation	Apply as full cover spray before flowering and repeat only when you have consulted with your agronomist or horticulturist.
Tomatoes	2 – 3 lbs/acre through spraying in 80 – 100 gal of water or 4 – 5 lbs/acre through irrigation	Apply as a full cover spray and repeat only when you have consulted with your agronomist or horticulturist.

AGROBOR™

F002806

GUARANTEED ANALYSIS

Boron (B)	16.00%
-----------	--------

Derived From: Boric Acid



Agri Technovation Inc.
516 Villa Ave, Ste 23,
Clovis, CA, 93612

**WATER-SOLUBLE CRYSTALLINE BORON COMPOUND**

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

KEEP OUT OF REACH OF CHILDREN.**DANGER** May damage fertility or the unborn child.

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Wear protective gloves, protective clothing and eye protection. IF exposed or concerned: Get medical advice. Store locked up. Dispose of container to in accordance with regulations.

NET WEIGHT: 50 LBS (22.68 kg)LIFT
HERE