

GUARANTEED ANALYSIS

Boron (B)	7.00 %
Molybdenum (Mo)	0.65 %

Derived From: Boron Ethanolamine, Sodium Molybdate.
ALSO INCLUDES NON PLANT FOOD INGREDIENT:
 20% Kelp (*Ascophyllum nodosum*, *Durvillaea potatorum*, *Ecklonia maxima*) (microbe food)



Agri Technovation Inc.
 516 Villa Ave, Ste 23,
 Clovis, CA, 93612



DENSITY: 10.26 LBS/GAL @ 68°F (20°C)

STORE ABOVE 40°F

LIQUID BORON AND MOLYBDENUM WITH ADDED KELP.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

KEEP OUT OF REACH OF CHILDREN

This product does not meet the hazardous criteria set forth by the 2012 OSHA Hazard Communication Standard (29 CFR 1910.1200). However, observe good industrial hygiene practices. Do not breathe spray. Wash face, hands and skin thoroughly after handling. Wear protective gloves, protective clothing and eye protection. Wash contaminated clothing before reuse. IF SWALLOWED: Rinse mouth. IF ON SKIN(or hair): Immediately remove all contaminated clothing. Rinse skin with water. IF INHALED: If breathing is difficult, move to fresh air and rest in a position comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

NET CONTENT: 265 GALLONS (1003.1 LITERS)

NET WEIGHT: 2 718.9 LBS (1233.27 KG)

DIRECTIONS FOR USE

KEL-MOB™ is recommended for use on broadacre and vegetable crops during plant growth periods that require relatively a high level of boron and molybdenum. KEL-MOB™ provides readily available nutrients as shown in the guaranteed analysis.

KEL-MOB™ will be influenced by differences in varieties, soil, and environmental conditions and should be applied under conditions of optimum plant growth. Optimum conditions are defined as adequate humidity, moderate temperatures, and adequate soil moisture.

- Wear suitable protective clothing when applied with other products.
- Optimum results with this product are obtained if the pH of the tank mix is between 5 and 6.
- For optimum results KEL-MOB™ should be used in a plant nutrition program. Recommendations should be based on soil and leaf analyses.
- Where overhead irrigation systems are in use, avoid scheduling irrigation directly after application.
- In orchards equipped with micro irrigation systems, spray deposits should not be affected by irrigation

CAUTION

Do not store mixed product for more than 8 hours. Consult with your distributor, crop adviser or manufacturer should crop specific programs, or any other information be required around the use of this product. Do not exceed the appropriate application rates.

Caution - KEL-MOB™ contains boron. Use on any crops other than those recommended on the label, may result in serious injury to the other crop(s).

Caution - KEL-MOB™ also contains molybdenum. Application of fertilizing materials containing molybdenum may result in forage crops containing levels of molybdenum which are toxic to ruminant animals.

STORAGE & DISPOSAL

Store product in original container. DO NOT store in excessive cold or heat or direct sunlight. DO NOT re-use container. DO NOT contaminate water, food or feed by storage or disposal. Keep packaging closed, securely fastened and upright. Triple rinse or equivalent and dispose of container in accordance with local, state, and federal regulations.

MIXING

Agitate KEL-MOB™ well before dilution or use. Always mix KEL-MOB™ with sufficient quantities of water. Never apply KEL-MOB™ undiluted. Follow standard tank mixing procedures. Ensure good agitation throughout application.

APPLICATION RATES

The rate of application is dependent on the soil fertility and reaction required. See table of recommended application rates for foliar application. For other modes of application, consult a qualified advisor. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies.

COMPATIBILITY

It is advisable to do a miscibility test prior to mixing with any other product and to consult with a qualified advisor or distributor.

DISCLAIMER

The manufacturer, importer and seller make no express or implied warranties, including any warranty of merchantability, fitness for a particular purpose, or suitability. The product is sold "AS-IS." All sales are final. In no event will manufacturer, importer and seller have any liability for any damages, whether direct, incidental, special or consequential arising out of or in connection with any duty with respect to the product, including, but not limited to lost profits, lost sales or injury to persons or property. Manufacturer, importer and seller's liability on any claim of any kind, including but not limited to warranty, negligence, strict liability, and any other cause of action, for any loss or damage arising out of, connected with, or resulting from the performance or breach of the terms of sale of any goods, or any related services related to sale, delivery, resale, technical direction, inspection, operation or use of any goods or services or part of those goods or services will in no case exceed the purchase price allocable to the goods or services or part of those goods or services that gives rise to the claim. Manufacturer, importer and seller make no representation whatsoever with regard to whether any goods comply with the Occupational Safety and Health Act (OSHA), the Americans with Disabilities Act (ADA), or any other federal, state or local statute, law, ordinance or ruling. Customer, buyer, and user assume all responsibility for use of product.

APPLICATION RATES

Crop	Maximum Rate per Acre	Rate per 100 gal water	Additional information
Barley	3.5 pt	8 pt	Spray one week before flowering.
Cabbage	3.5 pt	8 pt	Apply 2 – 3 weeks after plant and repeat the application 3 times at 21 day intervals.
Canola	3.5 pt	8 pt	Spray one week before flowering.
Carrots	3.5 pt	8 pt	Apply 2-3 weeks after 100% emergence and repeat 2-3 times at 21-day intervals.
Cauliflower	3.5 pt	8 pt	Apply 2-3 weeks after planting and repeat 3 times at 21-day intervals.
Cotton	3.5 pt	8 pt	Apply at the start of flowering and repeat in 14 days.
Peanuts	3.5 pt	8 pt	Apply 3 times at 21-day intervals starting 3 weeks after emergence.
Corn	4.5 pt	6 – 8 pt	Apply at V4 to V6 stage and repeat at silk emergence.

APPLICATION RATES

Crop	Maximum Rate per Acre	Rate per 100 gal water	Additional information
Soybeans	3.5 pt	16 pt	Application should take place between the V4 and V6 stage or just before rows close.
Drybeans	3.5 pt	16 pt	Application should take place between the V4 and V6 stage or just before rows close.
Sunflower	3.5 pt	16 pt	Spray one week before flowering.
Alfalfa	3.5 pt	16 pt	Spray at 10% flower and repeat if necessary.
Sorghum	3.5 pt	16 pt	Spray at 10% flower and repeat if necessary.
Potatoes	3.5 pt	16 pt	Apply 2 weeks after 100% emergence.
Tomatoes	3.5 pt	16 pt	Apply 2-3 weeks after 100% emergence and repeat 2-3 times at 21-day intervals.
Wheat	3.5 pt	16 pt	Spray one week before flowering.
Leafy vegetables	3.5 pt	16 pt	Apply 2-3 weeks after 100% emergence and repeat 2-3 times at 21-day intervals.

GUARANTEED ANALYSIS

Boron (B)	7.00 %
Molybdenum (Mo)	0.65 %

Derived From: Boron Ethanolamine, Sodium Molybdate.
ALSO INCLUDES NON PLANT FOOD INGREDIENT:
 20% Kelp (*Ascophyllum nodosum*, *Durvillaea potatorum*, *Ecklonia maxima*) (microbe food)



Agri Technovation Inc.
 516 Villa Ave, Ste 23,
 Clovis, CA, 93612



DENSITY: 10.26 LBS/GAL @ 68°F (20°C)

STORE ABOVE 40°F

LIQUID BORON AND MOLYBDENUM WITH ADDED KELP.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

KEEP OUT OF REACH OF CHILDREN

This product does not meet the hazardous criteria set forth by the 2012 OSHA Hazard Communication Standard (29 CFR 1910.1200). However, observe good industrial hygiene practices. Do not breathe spray. Wash face, hands and skin thoroughly after handling. Wear protective gloves, protective clothing and eye protection. Wash contaminated clothing before reuse. IF SWALLOWED: Rinse mouth. IF ON SKIN(or hair): Immediately remove all contaminated clothing. Rinse skin with water. IF INHALED: If breathing is difficult, move to fresh air and rest in a position comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

NET CONTENT: 265 GALLONS (1003.1 LITERS)

NET WEIGHT: 2 718.9 LBS (1233.27 KG)