

2 - 10 - 1

GUARANTEED ANALYSIS

| | |
|--|--------|
| Total Nitrogen (N) | 2.00% |
| 1.5% Ammoniacal Nitrogen | |
| 0.5% Nitrate Nitrogen | |
| Available Phosphate (P ₂ O ₅) | 10.00% |
| Soluble Potash (K ₂ O) | 1.00% |
| Magnesium (Mg) | 0.70% |
| 0.70% Water Soluble Magnesium (Mg) | |
| Boron (B) | 0.10% |
| Copper (Cu) | 0.10% |
| 0.10% Water Soluble Copper (Cu) | |
| Manganese (Mn) | 0.60% |
| 0.60% Water Soluble Manganese (Mn) | |
| Molybdenum (Mo) | 0.01% |
| Zinc (Zn) | 1.00% |
| 1.00% Water Soluble Zinc (Zn) | |



Agri Technovation Inc.

516 Villa Ave, Ste 23,
Clovis, CA, 93612



Derived From: Copper Sulfate, Kelp extract (*Ascophyllum nodosum*, *Durvillaea potatorum*, *Ecklonia maxima*), Magnesium Nitrate, Manganese Sulfate, Monoammonium Phosphate, Disodium Octaborate, Phosphoric Acid, Potassium Hydroxide, Sodium Molybdate, Zinc Sulfate

FOLIAR SPRAY TO PROMOTE GROWTH IN FRUIT CROPS. STORE ABOVE 40°F

DENSITY: 10.68 LBS/GAL @ 68°F (20°C)

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

KEEP OUT OF REACH OF CHILDREN.

DANGER Causes severe skin burns and eye damage. May damage fertility or the unborn child. May cause damage to organs through prolonged or repeated exposure.

Obtain special instructions before use. Do not handle until all the safety precautions have been read and understood. Do not breathe fumes. Wash face, hands and skin thoroughly after handling. Wear protective gloves, protective clothing and eye protection. IF SWALLOWED: Rinse mouth. Do not induce vomiting. IF ON SKIN (or hair): Immediately remove all contaminated clothing. Rinse skin with water. Wash clothing before reuse. IF INHALED: If breathing is difficult, remove victim to fresh air and keep at rest in a position comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If exposed or concerned: Immediately call a POISON CENTER or doctor. Store locked up. Dispose of container in accordance with regulations. 14.3% of ingredients in mixture has unknown acute toxicity.

NET CONTENTS: 265 GALLONS (1003 Liters)

NET WEIGHT: 2830 (1283 kg)

DIRECTIONS FOR USE

MAJESTIC™ is recommended for use during plant growth periods that require relatively high levels of macro and micro nutrients. MAJESTIC™ is an effective, readily available source of the nutrients shown in the guaranteed analysis for foliar applications.

MAJESTIC™ will be influenced by differences in crop varieties, soil and environmental conditions and should be applied under conditions of optimum plant growth. Optimum conditions are defined as adequate humidity, moderate temperatures and adequate soil moisture. Optimum conditions for foliar application to most crops generally occur during early morning or late evening hours.

- Do not apply to crops that are water or heat stressed or in very windy conditions.
- Wear suitable protective clothing when sprayed with other products.
- For best results with this product, the pH of the tank mixture should be between 5 and 6.
- This product should be applied as a foliar spray when crops have sufficient leaf area.
- For optimum results MAJESTIC™ should be used in a plant nutrition program. Recommendations should be based on soil and leaf analyses.
- When applied as a soil application or fertigation do not exceed 1 gal per acre.
- Use a lower dosage per 100 gal at higher water volumes and a higher dosage per 100 gal at lower spray volumes.
- Where overhead systems are in use, avoid scheduling irrigation directly after application.
- In orchards equipped with micro irrigation systems, spray deposits should not be affected by irrigation.

CAUTION

Do not apply on wet fruit. Application can only take place once the fruit is dry. Do not apply to plants that are undergoing a period of moisture or heat stress, or during cool wet conditions. Do not spray MAJESTIC™ within 30 days after application of any Copper containing products. Do not store mixed product for more than 8 hours. Do NOT mix or apply in combination with Calcium containing products, amine herbicides, Copper containing fungicides, Sulfates, Phosphates, phosphonates, highly alkaline material such as Lime Sulfur, MCPA or highly alkaline Boron products. Do not exceed the appropriate application rates.

CAUTION

MAJESTIC™ contains Boron. Use of MAJESTIC™ on any crops other than those recommended may result in serious injury to the crop(s).

CAUTION

MAJESTIC™ also contains Molybdenum. Application of fertilizing materials containing Molybdenum may result in forage crops containing levels of Molybdenum which are toxic to ruminant animals.

Consult with your distributor, crop adviser or manufacturer should crop specific programs, or any other information be required around the use of this product.

STORAGE & DISPOSAL

Store product in original container. DO NOT store in excessive cold or heat or direct sunlight. DO NOT reuse container. DO NOT contaminate water, food or feed by storage or disposal. Keep packaging closed, securely fastened and upright. Triple rinse or equivalent and dispose of container in accordance with local, state and federal regulations.

MIXING

Agitate MAJESTIC™ well before dilution or use. Always mix MAJESTIC™ with sufficient quantities of water. Never apply MAJESTIC™ undiluted. Follow standard tank mixing procedures. Ensure good agitation throughout application.

COMPATIBILITY

It is advisable to do a miscibility test prior to mixing with any other product and to consult with a qualified advisor or distributor.

APPLICATION RATES

The rate of application is dependent on the soil fertility and reaction required. For more information on foliar application, see table on application rates. For other modes of application, consult a qualified advisor. Lower rates should be used for maintenance while higher rates will correct nutritional deficiencies.

DISCLAIMER

The manufacturer, importer and seller make no express or implied warranties, including any warranty of merchantability, fitness for a particular purpose, or suitability. The product is sold "AS-IS". All sales are final. In no event will manufacturer, importer and seller have any liability for any damages, whether direct, incidental, special or consequential arising out of or in connection with any duty with respect to the product, including, but not limited to lost profits, lost sales or injury to persons or property. Manufacturer, importer and seller's liability on any claim of any kind, including but not limited to warranty, negligence, strict liability, and any other cause of action, for any loss or damage arising out of, connected with, or resulting from the performance or breach of the terms of sale of any goods, or any related services related to sale, delivery, resale, technical direction, inspection, operation or use of any goods or services or part of those goods or services will in no case exceed the purchase price allocable to the goods or services or part of those goods or services that gives rise to the claim. Manufacturer, importer and seller make no representation whatsoever with regard to whether any goods comply with the Occupational Safety and Health Act (OSHA), the Americans with Disabilities Act (ADA), or any other federal, state or local statute, law, ordinance or ruling. Customer, buyer and user assume all responsibility for use of product.

FOLIAR APPLICATION RATES

| Crop | Max rate per acre | Rate per 100 gal Water | Additional Information |
|----------------------------|-------------------|------------------------|---|
| Pecan nuts | 2 - 3 qt | 1.5 pt | Apply after leaves have emerged and when enough leaves have developed for nutrient uptake. Apply once every two weeks for the first two months. |
| Macadamias | 2 - 3 qt | 1.5 pt | First application at pre-blossom. Second application at 100% petal drop and third application, four weeks after 100% petal drop. |
| Bearing apples and pears | 2 - 3 qt | 1.5 pt | First application after leaves have emerged and when enough leaves have developed for nutrient uptake. Second application 14 days later. |
| Citrus | 2 - 3 qt | 1.5 pt | First application at pre-blossom. Second application at 100% petal drop and third application, four weeks after 100% petal drop. Fourth application at flower initiation. |
| Avocados, mangoes, lychees | 2 - 3 qt | 1.5 pt | First application at pre-blossom. Second application at 100% petal drop and third application, four weeks after 100% petal drop. |
| Olives | 2 - 3 qt | 1.5 pt | First application at pre-blossom. Second application at 100% petal drop and third application, four weeks after 100% petal drop. Fourth application at flower initiation. |
| Stone fruit | 2 - 3 qt | 1.5 pt | First application after leaves have emerged and when enough leaves have developed for nutrient uptake. Second application 14 days later. |

FOLIAR APPLICATION RATES

| Crop | Max rate per acre | Rate per 100 gal Water | Additional Information |
|---|-------------------|------------------------|--|
| Table grapes | 1 - 2 qt | 2 pt | First application when shoots are 4 inches. Further applications at end of flowering and when berries are pea size. Apply one application at post-harvest. |
| Wine grapes | 1 - 2 qt | 1.5 pt | First application when shoots are 4 inches, repeat spray 2 weeks later. |
| Bananas | 2 - 3 qt | 1.5 pt | Start spraying when suckers reach 40 inches in length, repeat twice, fortnightly. |
| Blueberries, strawberries and gooseberries. | 2 qt | 1.5 pt | First application at pre-blossom. Second at 100% petal drop. |
| Almonds | 2 - 3 qt | 1.5 pt | First application at pre-blossom. Second one two weeks after bloom and third application 5 weeks after bloom. |
| Pistachios | 2 - 3 qt | 1.5 pt | First application at pre-blossom. Second application 10 days after bloom and third application 25 days after bloom. |

2 - 10 - 1

GUARANTEED ANALYSIS

| | |
|--|--------|
| Total Nitrogen (N) | 2.00% |
| 1.5% Ammoniacal Nitrogen | |
| 0.5% Nitrate Nitrogen | |
| Available Phosphate (P ₂ O ₅) | 10.00% |
| Soluble Potash (K ₂ O) | 1.00% |
| Magnesium (Mg) | 0.70% |
| 0.70% Water Soluble Magnesium (Mg) | |
| Boron (B) | 0.10% |
| Copper (Cu) | 0.10% |
| 0.10% Water Soluble Copper (Cu) | |
| Manganese (Mn) | 0.60% |
| 0.60% Water Soluble Manganese (Mn) | |
| Molybdenum (Mo) | 0.01% |
| Zinc (Zn) | 1.00% |
| 1.00% Water Soluble Zinc (Zn) | |



Agri Technovation Inc.

516 Villa Ave, Ste 23,
Clovis, CA, 93612



Derived From: Copper Sulfate, Kelp extract (*Ascophyllum nodosum*, *Durvillaea potatorum*, *Ecklonia maxima*), Magnesium Nitrate, Manganese Sulfate, Monoammonium Phosphate, Disodium Octaborate, Phosphoric Acid, Potassium Hydroxide, Sodium Molybdate, Zinc Sulfate

FOLIAR SPRAY TO PROMOTE GROWTH IN FRUIT CROPS. STORE ABOVE 40°F

DENSITY: 10.68 LBS/GAL @ 68°F (20°C)

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

KEEP OUT OF REACH OF CHILDREN.

DANGER Causes severe skin burns and eye damage. May damage fertility or the unborn child. May cause damage to organs through prolonged or repeated exposure.

Obtain special instructions before use. Do not handle until all the safety precautions have been read and understood. Do not breathe fumes. Wash face, hands and skin thoroughly after handling. Wear protective gloves, protective clothing and eye protection. IF SWALLOWED: Rinse mouth. Do not induce vomiting. IF ON SKIN (or hair): Immediately remove all contaminated clothing. Rinse skin with water. Wash clothing before reuse. IF INHALED: If breathing is difficult, remove victim to fresh air and keep at rest in a position comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If exposed or concerned: Immediately call a POISON CENTER or doctor. Store locked up. Dispose of container in accordance with regulations. 14.3% of ingredients in mixture has unknown acute toxicity.

NET CONTENTS: 265 GALLONS (1003 Liters)

NET WEIGHT: 2830 (1283 kg)